

Ordering and Billing EXPAREL for Your Facility

» A Proven, Long-Lasting, Non-Opioid Option

EXPAREL is indicated for single-dose infiltration in patients aged 6 years and older to produce postsurgical local analgesia and in adults as an interscalene brachial plexus nerve block to produce postsurgical regional analgesia. Safety and efficacy have not been established in other nerve blocks.

NDC # (on package label)	Dose/Vial Size	Configuration	NDC # (for billing)*
65250-133-10	EXPAREL 133 mg/10 mL	Single-dose vial	65250-0133-10
65250-266-20	EXPAREL 266 mg/20 mL	Single-dose vial	65250-0266-20



EXPAREL is available in 2 doses/vial sizes: 133 mg/10 mL and 266 mg/20 mL.

» Product Configurations

Both doses of EXPAREL can be purchased in cartons of 4 or 10 vials. Consult the information below when ordering or billing for EXPAREL.

NDC # (on package label)	Dose/Vial Size	Configuration	NDC # (for billing)*
65250-133-04	EXPAREL 133 mg/10 mL	Carton of 4 vials	65250-0133-10
65250-133-09	EXPAREL 133 mg/10 mL	Carton of 10 vials	65250-0133-10
65250-266-04	EXPAREL 266 mg/20 mL	Carton of 4 vials	65250-0266-20
65250-266-09	EXPAREL 266 mg/20 mL	Carton of 10 vials	65250-0266-20



Two Ways to Order

Wholesaler

Contact your wholesaler to order EXPAREL today!

OR

Pacira BioSciences, Inc.

Representative: _____
 Email: _____
 Phone: _____

NDC=National Drug Code.

*10-digit NDC numbers should always be converted to the appropriate 11-digit NDC number when billing for reimbursement.

Please see Important Safety Information on reverse.
 Please refer to accompanying full Prescribing Information, which is also available at www.EXPAREL.com.

NON-OPIOID
EXPAREL®
 (bupivacaine liposome injectable suspension)

**NON-OPIOID
EXPAREL®**
(bupivacaine liposome injectable suspension)

Indication

EXPAREL is indicated for single-dose infiltration in patients aged 6 years and older to produce postsurgical local analgesia and in adults as an interscalene brachial plexus nerve block to produce postsurgical regional analgesia. Safety and efficacy have not been established in other nerve blocks.

Important Safety Information

EXPAREL is contraindicated in obstetrical paracervical block anesthesia.

Adverse reactions reported in adults with an incidence greater than or equal to 10% following EXPAREL administration via infiltration were nausea, constipation, and vomiting; adverse reactions reported in adults with an incidence greater than or equal to 10% following EXPAREL administration via interscalene brachial plexus nerve block were nausea, pyrexia, and constipation.

Adverse reactions with an incidence greater than or equal to 10% following EXPAREL administration via infiltration in pediatric patients six to less than 17 years of age were nausea, vomiting, constipation, hypotension, anemia, muscle twitching, vision blurred, pruritus, and tachycardia.

If EXPAREL and other non-bupivacaine local anesthetics, including lidocaine, are administered at the same site, there may be an immediate release of bupivacaine from EXPAREL. Therefore, EXPAREL may be administered to the same site 20 minutes after injecting lidocaine. EXPAREL is not recommended to be used in the following patient populations: patients <6 years old for infiltration, patients younger than 18 years old for interscalene brachial plexus nerve block, and/or pregnant patients.

Because amide-type local anesthetics, such as bupivacaine, are metabolized by the liver, EXPAREL should be used cautiously in patients with hepatic disease.

Warnings and Precautions Specific to EXPAREL

Avoid additional use of local anesthetics within 96 hours following administration of EXPAREL.

EXPAREL is not recommended for the following types or routes of administration: epidural, intrathecal, regional nerve blocks **other than interscalene brachial plexus nerve block**, or intravascular or intra-articular use.

The potential sensory and/or motor loss with EXPAREL is temporary and varies in degree and duration depending on the site of injection and dosage administered and may last for up to 5 days, as seen in clinical trials.

Warnings and Precautions for Bupivacaine-Containing Products

Central Nervous System (CNS) Reactions: There have been reports of adverse neurologic reactions with the use of local anesthetics. These include persistent anesthesia and paresthesia. CNS reactions are characterized by excitation and/or depression.

Cardiovascular System Reactions: Toxic blood concentrations depress cardiac conductivity and excitability which may lead to dysrhythmias, sometimes leading to death.

Allergic Reactions: Allergic-type reactions (eg, anaphylaxis and angioedema) are rare and may occur as a result of hypersensitivity to the local anesthetic or to other formulation ingredients.

Chondrolysis: There have been reports of chondrolysis (mostly in the shoulder joint) following intra-articular infusion of local anesthetics, which is an unapproved use.

Methemoglobinemia: Cases of methemoglobinemia have been reported with local anesthetic use.

Please refer to accompanying full Prescribing Information, which is also available at www.EXPAREL.com.

For more information, please visit www.EXPAREL.com or call 1-855-792-9727.